

INSTRUCTIONS: Circle the number that applies to you. **If a symptom does not apply, leave it blank.**
 Circle either: (1) for **MILD** symptoms (occurs rarely), (2) for **MODERATE** symptoms (occurs several times a month),
 or (3) for **SEVERE** symptoms (occurs almost constantly).

GROUP ONE

- | | | |
|-----------------------------------|--|-----------------------------------|
| 1 - 1 2 3 Acid foods upset | 8 - 1 2 3 Gag Easily | 15 - 1 2 3 Appetite reduced |
| 2 - 1 2 3 Get chilled, often | 9 - 1 2 3 Unable to relax, startles easily | 16 - 1 2 3 Cold sweats often |
| 3 - 1 2 3 "Lump" in throat | 10 - 1 2 3 Extremities cold, clammy | 17 - 1 2 3 Fever easily raised |
| 4 - 1 2 3 Dry mouth-eyes-nose | 11 - 1 2 3 Strong light irritates | 18 - 1 2 3 Neuralgia-like pains |
| 5 - 1 2 3 Pulse speeds after meal | 12 - 1 2 3 Urine amount reduced | 19 - 1 2 3 Staring, blinks little |
| 6 - 1 2 3 Keyed up - fail to calm | 13 - 1 2 3 Heart pounds after retiring | 20 - 1 2 3 Sour stomach frequent |
| 7 - 1 2 3 Cuts heal slowly | 14 - 1 2 3 "Nervous" stomach | |

GROUP TWO

- | | | |
|---|--|---------------------------------------|
| 21 - 1 2 3 Joint stiffness after arising | 29 - 1 2 3 Digestion rapid | 37 - 1 2 3 "Slow starter" |
| 22 - 1 2 3 Muscle-leg-toe cramps at night | 30 - 1 2 3 Vomiting frequent | 38 - 1 2 3 Get "chilled" infrequently |
| 23 - 1 2 3 "Butterfly" stomach, cramps | 31 - 1 2 3 Hoarseness frequent | 39 - 1 2 3 Perspire easily |
| 24 - 1 2 3 Eyes or nose watery | 32 - 1 2 3 Breathing irregular | 40 - 1 2 3 Circulation poor, |
| 25 - 1 2 3 Eyes blink often | 33 - 1 2 3 Pulse slow; feels "irregular" | sensitive to cold |
| 26 - 1 2 3 Eyelids swollen, puffy | 34 - 1 2 3 Gagging reflex slow | 41 - 1 2 3 Subject to colds, |
| 27 - 1 2 3 Indigestion soon after meals | 35 - 1 2 3 Difficulty swallowing | asthma, bronchitis |
| 28 - 1 2 3 Always seem hungry;
feels "lightheaded" often | 36 - 1 2 3 Constipation,
diarrhea alternating | |

GROUP THREE

- | | | |
|---|--|---|
| 42 - 1 2 3 Eat when nervous | 49 - 1 2 3 Heart palpitates if meals
missed or delayed | 53 - 1 2 3 Crave candy or coffee
in afternoons |
| 43 - 1 2 3 Excessive appetite | 50 - 1 2 3 Afternoon headaches | 54 - 1 2 3 Moods of depression -
"blues" or melancholy |
| 44 - 1 2 3 Hungry between meals | 51 - 1 2 3 Overeating sweets upsets | 55 - 1 2 3 Abnormal craving for
sweets or snacks |
| 45 - 1 2 3 Irritable before meals | 52 - 1 2 3 Awaken after few hours sleep
- hard to get back to sleep | |
| 46 - 1 2 3 Get "shaky" if hungry | | |
| 47 - 1 2 3 Fatigue, eating relieves | | |
| 48 - 1 2 3 "Lightheaded" if meals delayed | | |

GROUP FOUR

- | | | |
|---|---|--|
| 56 - 1 2 3 Hands and feet go to sleep
easily, numbness | 63 - 1 2 3 Get "drowsy" often | 68 - 1 2 3 Bruise easily, "black
and blue" spots |
| 57 - 1 2 3 Sigh frequently, "air
hunger" | 64 - 1 2 3 Swollen ankles
worse at night | 69 - 1 2 3 Tendency to anemia |
| 58 - 1 2 3 Aware of "breathing
heavily" | 65 - 1 2 3 Muscle cramps, worse
during exercise; get
"charley horses" | 70 - 1 2 3 "Nose bleeds" frequent |
| 59 - 1 2 3 High altitude discomfort | 66 - 1 2 3 Shortness of breath
on exertion | 71 - 1 2 3 Noises in head, or
"ringing in ears" |
| 60 - 1 2 3 Opens windows in
closed room | 67 - 1 2 3 Dull pain in chest or
radiating into left arm,
worse on exertion | 72 - 1 2 3 Tension under the
breastbone, or feeling
of "tightness",
worse on exertion |
| 61 - 1 2 3 Susceptible to colds
and fevers | | |
| 62 - 1 2 3 Afternoon "yawner" | | |